

A Green's function formalism of energy and momentum transfer in fluctuational electrodynamics

A. Narayanaswamy^{a,*}, Y. Zheng^a

^aDepartment of Mechanical Engineering, Columbia University, New York, NY 10027

Abstract

Radiative energy and momentum transfer due to fluctuations of electromagnetic fields arising due to temperature difference between objects is described in terms of the cross-spectral densities of the electromagnetic fields. We derive relations between thermal non-equilibrium contributions to energy and momentum transfer and surface integrals of tangential components of the dyadic Green's functions of the vector Helmholtz equation. The expressions derived here are applicable to objects of arbitrary shapes, dielectric functions, as well as magnetic permeabilities. For the case of radiative transfer, we derive expressions for the generalized transmissivity and generalized conductance that are shown to obey reciprocity and agree with theory of black body radiative transfer in the appropriate limit.

Keywords: Dyadic Green's function, Near field radiative transfer, Fluctuational electrodynamics, Non-equilibrium effects

PACS: 41.20.-q, 41.20.Jb, 42.25.Bs

Nomenclature

$\mathbf{E}$	Electric field vector	T^e	Generalized transmissivity for radiative energy transfer
F	View factor	T^m	Generalized transmissivity for momentum transfer
$\overline{\overline{G}}_e$	Electric dyadic Green's function	V_l	Volume of object l
$\overline{\overline{G}}_m$	Magnetic dyadic Green's function	V_δ	Volume of infinitesimal radius surrounding $\tilde{\mathbf{r}}$
$\overline{\overline{G}}_E$	$\nabla \times \overline{\overline{G}}_e$	$\overline{\overline{\mathcal{E}}}$	Matrix of contribution to $\langle \mathbf{EE}^* \rangle_s$
$\overline{\overline{G}}_M$	$\nabla \times \overline{\overline{G}}_m$	$\overline{\overline{\mathcal{H}}}$	Matrix of contribution to $\langle \mathbf{HH}^* \rangle_s$
$\overline{\overline{G}}_o$	Green's function of contribution due to background or source radiation	$\overline{\overline{\mathcal{X}}}$	Matrix of contribution to $\langle \mathbf{EH}^* \rangle_s$
$\overline{\overline{G}}^{(sc)}$	Green's function of contribution from waves scattered by interfaces	c	Speed of light
G^e	Linearized conductance for radiative transfer	$\hbar$	Reduced Planck's constant
$\mathbf{H}$	Magnetic field vector	k	Wavevector
$\overline{\overline{I}}$	Identity matrix	k_b	Boltzmann's constant
$\mathbf{J}$	Current Density	k_n	n component of wavevector ($n = x, y, z$)
T_l	Temperature in object l	k_{hz}	z component of wavevector in vacuum
$\mathbf{P}$	Poynting vector	k_ρ	$\sqrt{k_x^2 + k_y^2}$
Q	Radiative heat transfer	l	Thickness of vacuum gap
$\tilde{R}_{hn}$	Fresnel reflection coefficients at interfaces between h and n	$\hat{\mathbf{n}}$	Unit normal vector
S_l	Closed surface of object l	$\mathbf{r}$	Position vector of observation point
		$\tilde{\mathbf{r}}$	Position vector of source point
		t	Time
		β	0 or 1
		δ	Delta function
		ρ	Distance of points
		ϵ	Levi-Civita symbol

*Corresponding author

Email address: arvind.narayanaswamy@columbia.edu (A. Narayanaswamy)

ε	Permittivity, $\varepsilon' + i\varepsilon''$
ε_o	Permittivity of free space
Θ	Energy of a photon at temperature T
μ	Permeability, $\mu' + i\mu''$
μ_o	Permeability of free space
ν, ξ	1 or -1
$\overline{\overline{\sigma}}$	Maxwell stress tensor
ω	Frequency
$\Re$	Real part
$\Im$	Imaginary part
Tr	Trace

Superscripts

bb	Blackbody
e	Electric field
m	Magnetic field
pp	Planar-planar
(h)	Vacuum
(l)	Objects ($l = 1, 2, \dots, N$)
(p)	Transverse magnetic
(s)	Transverse electric
(μ)	Polarization s or p
T	Transpose
$*$	Complex conjugate

Subscripts

h	Vacuum
i, p, q	Cartesian components 1, 2, 3
l	Objects ($l = 1, 2, \dots, N$)
s	Symmetric summation
$1 \rightarrow 2$	From object 1 to 2

1. Introduction

Fluctuations of electromagnetic fields lead to thermal radiative transfer, via energy transfer, and van der Waals and Casimir forces, via momentum transfer. Diffraction and interference effects as well as tunneling of evanescent and surface waves, collectively known as near-field effects, are not taken into consideration by the classical theory of radiative transfer. Near-field effects become important when the length scale of importance becomes comparable to the characteristic thermal wavelength ($\lambda_T \approx 3000/T$ μm). For radiative transfer between two objects, an important length scale is the minimum inter-object spacing, l_{gap} . When $l_{gap} \ll \lambda_T$, tunneling of electromagnetic waves lead to enhancement of radiative transfer beyond the classical or far-field limit. Surface texturing, for instance by creating a periodic 1D or 2D pattern, introduces a length scale,

l_p , that characterizes the period of the pattern. When $l_p \ll \lambda_T$, diffraction effects can lead to thermal emission patterns not usually associated with a planar surface [1].

It has been long recognized that near-field enhancement of radiative transfer due to surface polaritons can result in increased power density as well as efficiency [2, 3, 4, 5]. However, this enhancement of energy transfer has not been used in any practical device, as yet, because of our inability to conceive of configurations other than two parallel surfaces with a thin vacuum gap in which an enhancement of similar magnitude occurs. Most investigations of near-field radiative transfer have been restricted to objects of few simple geometric shapes, each analyzed by a vector eigenfunction expansion method applicable to that geometry (planar geometry with vector plane waves [2, 6, 7, 8, 9, 10], cylindrical surfaces with vector cylindrical waves [11], two spheres with vector spherical waves [12, 13, 14, 15], sphere-plane with a combination of vector spherical and plane waves [16]). Even minor changes to the shape of the object can impose great challenges. Simulations of thermal emission from textured surfaces are usually performed using rigorous coupled wave analysis (RCWA) [17, 18, 19] or finite difference time domain (FDTD) methods [20], which are quite different from those used for simulations of near-field radiative transfer. To design other types of surfaces that can exploit the enhancement, without posing the hurdles associated with two parallel surfaces, and also to design surfaces with new radiative properties by shape modification at nano/micro scale, we need a general method to predict all types of nanoscale effects on radiative transfer, irrespective of the size, shape or properties of the objects involved.

Kruger et al. [11, 21] used fluctuational electrodynamics to develop a scattering matrix and operator formalism for computing non-equilibrium force and heat transfer interactions between objects with arbitrary shapes and frequency dependent dielectric permittivities. Biehs et al. [9] developed a formalism of nanoscale radiative transfer between two parallel surfaces similar to that of Landauer formalism of electron transport in mesoscopic devices [22, 23, 24, 25]. Ben-Abdallah et al. [26] used Rytov's theory to develop a theoretical formalism for radiative transfer between many objects in the dipole limit. Messina et al. [27] proposed a scattering matrix version of nanoscale radiative transfer as well as dispersion forces that is valid for objects with arbitrary shapes as well as dielectric functions. Non-equilibrium fluctuational electrodynamical interactions between objects can be expressed in a scattering matrix formalism or in a Green's functions formalism, just as the electrical conductance for electron transport can be developed in terms of the scattering matrix or Green's function.

The work in this paper is an extension to a prior work published in this journal by one of the authors [28]. In Ref. [28], the focus was on the relation between cross-spectral densities of electromagnetic fields in thermal equilibrium and the dyadic Green's functions (DGFs) of the vector




Figure 1: Schematic of N objects at temperatures $T_1, T_2, \dots, T_N$ embedded in a host medium at T_h .

Helmholtz equation. In this paper, the focus is on thermal non-equilibrium effects, i.e. when the objects are at different temperatures. Volume integral expressions for cross-spectral densities of components of the electric and magnetic fields that can be obtained from Rytov's theory of fluctuational electrodynamics are converted into a form more appropriate (in terms of surface integrals of DGFs of the vector Helmholtz equation) for computations as well as comparison with the classical theory of radiative transfer. Though the focus of this paper is not on developing new numerical techniques, it is hoped that the formalism developed here will be used to compute thermal non-equilibrium energy and momentum transfer between arbitrarily shaped objects.

The paper is arranged as follows. In Sec. 2, the fluctuation-dissipation theorem and DGFs are used to express the electric and magnetic field correlation functions in terms of the volume integrals of the DGFs. In Sec. 3, Green's identities for dyadic functions are used to derive expressions for the field correlations in terms of surface integrals of tangential components of the DGFs. For radiative transfer between two objects, a generalized transmissivity function that is expressed in terms of double surface integrals on the surfaces of the two objects is derived in Sec. 4. In Sec. 5, the theoretical formalism developed in Sec. 4 is applied to the cases of radiative heat transfer between two parallel half spaces. We also show that the generalized transmissivity function agrees with the theory of blackbody radiation in the appropriate limit. Finally, we also discuss the implications of the theoretical formalism developed here for computation of heat transfer and non-equilibrium forces.

2. Fluctuational electrodynamics and Green's function formalism

We briefly describe our notation regarding electromagnetic fields and their Fourier transforms here. A field $A(\mathbf{r}, t)$ and its Fourier transform, $A(\mathbf{r}, \omega)$, are related by

$A(\mathbf{r}, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} A(\mathbf{r}, \omega) e^{-i\omega t} d\omega$. Since the same symbol is used to identify a field as well as its Fourier transform, explicit dependence on time will be included when referring to the time domain field. Explicit dependence on ω is suppressed from $A(\mathbf{r}, \omega)$ so that it is written as $A(\mathbf{r})$. Explicit dependence of relative dielectric permittivities, magnetic permeabilities, and DGFs on ω is also suppressed.

Let us consider N objects (see Fig. 1) with relative dielectric permittivities $\epsilon_l(\omega)$ and magnetic permeabilities $\mu_l(\omega)$ at temperatures T_l , where $l = 1, 2, \dots, N$. These objects are assumed to be embedded in vacuum that is at temperature T_h . The object l is confined to the volume V_l and the closed surface S_l is the boundary of this object with the host medium. The outward normal on the surface of object l at $\mathbf{r}$ is represented by $\mathbf{n}_l(\mathbf{r})$. The fluctuations of the electric and magnetic current densities, which give rise to the dispersion forces and radiative transfer, are related to temperature by fluctuation-dissipation theorems of the *second kind* [29, 30, 31]:

$$\langle J_p^e(\mathbf{r}) J_q^{e*}(\tilde{\mathbf{r}}) \rangle = 2\omega \epsilon_o \epsilon'' \Theta(\omega, T) \delta(\mathbf{r} - \tilde{\mathbf{r}}) \delta_{pq} \quad (1a)$$

$$\langle J_p^m(\mathbf{r}) J_q^{m*}(\tilde{\mathbf{r}}) \rangle = 2\omega \mu_o \mu'' \Theta(\omega, T) \delta(\mathbf{r} - \tilde{\mathbf{r}}) \delta_{pq} \quad (1b)$$

$$\langle J_p^e(\mathbf{r}) J_q^{m*}(\tilde{\mathbf{r}}) \rangle = 0 \quad (1c)$$

where $p, q = 1, 2, 3$ are the labels for the Cartesian components of the vector, ϵ_o and μ_o are the permittivity and permeability of free space, J_p^e and J_p^m are the Cartesian components of the electric and magnetic current densities, $\Theta(\omega, T) = \frac{\hbar\omega}{2} \coth\left(\frac{\hbar\omega}{2k_b T}\right)$, ϵ'' and μ'' are the imaginary parts of the dielectric permittivity and magnetic permeability respectively at the location $\mathbf{r}$ which is in local thermodynamic equilibrium at temperature T , z^* is the complex conjugate of z , and $\langle \rangle$ denotes the ensemble average. $2\pi\hbar$ is the Planck constant and k_b is the Boltzmann constant. The presence of δ_{pq} implies that we assume all materials to be isotropic, and that of $\delta(\mathbf{r} - \tilde{\mathbf{r}})$ implies that the correlations of sources are local. The Fourier transforms of the electric and magnetic fields in the host medium (volume V_h in Fig. 1) due to sources in object l (volume V_l) are given by:

$$\mathbf{E}(\tilde{\mathbf{r}}) = \int_{V_l} [\mathbf{p}(\mathbf{r}) \cdot \overline{\mathbf{G}}_e(\mathbf{r}, \tilde{\mathbf{r}}) - \mathbf{J}^m(\mathbf{r}) \cdot \overline{\mathbf{G}}_E(\mathbf{r}, \tilde{\mathbf{r}})] d\mathbf{r} \quad (2a)$$

$$\mathbf{H}(\tilde{\mathbf{r}}) = \int_{V_l} [\mathbf{m}(\mathbf{r}) \cdot \overline{\mathbf{G}}_m(\mathbf{r}, \tilde{\mathbf{r}}) + \mathbf{J}^e(\mathbf{r}) \cdot \overline{\mathbf{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})] d\mathbf{r} \quad (2b)$$

where $\mathbf{p}(\mathbf{r}) = i\omega\mu_o\mu(\mathbf{r})\mathbf{J}^e(\mathbf{r})$, $\mathbf{m}(\mathbf{r}) = i\omega\epsilon_o\epsilon(\mathbf{r})\mathbf{J}^m(\mathbf{r})$, $\tilde{\mathbf{r}} \in V_h$, $\overline{\mathbf{G}}_E(\mathbf{r}, \tilde{\mathbf{r}}) = \nabla \times \overline{\mathbf{G}}_e(\mathbf{r}, \tilde{\mathbf{r}})$ and $\overline{\mathbf{G}}_M(\mathbf{r}, \tilde{\mathbf{r}}) = \nabla \times \overline{\mathbf{G}}_m(\mathbf{r}, \tilde{\mathbf{r}})$. $\overline{\mathbf{G}}_e(\mathbf{r}, \tilde{\mathbf{r}})$ and $\overline{\mathbf{G}}_m(\mathbf{r}, \tilde{\mathbf{r}})$ are DGFs of the vector Helmholtz equation that satisfy the following boundary conditions on the interface S_l between object l and the host medium:

$$\hat{\mathbf{n}}_l(\mathbf{r}_l) \times (\mu_l(\mathbf{r}_l) \overline{\mathbf{G}}_e(\mathbf{r}_l, \tilde{\mathbf{r}}) - \mu_h(\mathbf{r}_h) \overline{\mathbf{G}}_e(\mathbf{r}_h, \tilde{\mathbf{r}})) = 0, \quad (3a)$$

$$\hat{\mathbf{n}}_l(\mathbf{r}_l) \times (\bar{\bar{\mathbf{G}}}_E(\mathbf{r}_l, \tilde{\mathbf{r}}) - \bar{\bar{\mathbf{G}}}_E(\mathbf{r}_h, \tilde{\mathbf{r}})) = 0, \quad (3b)$$

$$\hat{\mathbf{n}}_l(\mathbf{r}_l) \times (\varepsilon_l(\mathbf{r}_l)\bar{\bar{\mathbf{G}}}_m(\mathbf{r}_l, \tilde{\mathbf{r}}) - \varepsilon_h(\mathbf{r}_h)\bar{\bar{\mathbf{G}}}_m(\mathbf{r}_h, \tilde{\mathbf{r}})) = 0, \quad (3c)$$

$$\hat{\mathbf{n}}_l(\mathbf{r}_l) \times (\bar{\bar{\mathbf{G}}}_M(\mathbf{r}_l, \tilde{\mathbf{r}}) - \bar{\bar{\mathbf{G}}}_M(\mathbf{r}_h, \tilde{\mathbf{r}})) = 0, \quad (3d)$$

where $\mathbf{r}_l$ and $\mathbf{r}_h$ are position vectors of points on either side of S_l in volume V_l and V_h respectively ($|\mathbf{r}_l - \mathbf{r}_h| \rightarrow 0$). In addition, the DGFs satisfy the following reciprocity relations:

$$\mu(\mathbf{r})\bar{\bar{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) = \mu(\tilde{\mathbf{r}})\bar{\bar{G}}_e(\tilde{\mathbf{r}}, \mathbf{r}), \quad (4)$$

$$\varepsilon(\mathbf{r})\bar{\bar{G}}_m^T(\mathbf{r}, \tilde{\mathbf{r}}) = \varepsilon(\tilde{\mathbf{r}})\bar{\bar{G}}_m(\tilde{\mathbf{r}}, \mathbf{r}), \quad (5)$$

$$\bar{\bar{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) = \bar{\bar{G}}_M(\tilde{\mathbf{r}}, \mathbf{r}), \quad (6)$$

where $\bar{\bar{A}}^T$ is the transpose of $\bar{\bar{A}}$.

Radiative transfer can be determined from the Poynting vector, $\mathbf{P}(\tilde{\mathbf{r}}) = \langle \mathbf{E}(\tilde{\mathbf{r}}, t) \times \mathbf{H}(\tilde{\mathbf{r}}, t) \rangle$, whose components are given by:

$$P_i(\tilde{\mathbf{r}}) = \epsilon_{ipq} \langle E_p(\tilde{\mathbf{r}}, t) H_q(\tilde{\mathbf{r}}, t) \rangle \quad (7)$$

where ϵ_{ipq} is the Levi-Civita symbol. To determine van der Waals pressure and radiative transfer, we need equal time correlations of various components of the electric and magnetic field vectors, such as $\langle E_p(\tilde{\mathbf{r}}, t) E_q(\tilde{\mathbf{r}}, t) \rangle$, $\langle H_p(\tilde{\mathbf{r}}, t) H_q(\tilde{\mathbf{r}}, t) \rangle$, and $\langle E_p(\tilde{\mathbf{r}}, t) H_q(\tilde{\mathbf{r}}, t) \rangle$. van der Waals pressure in vacuum can be determined from the Maxwell stress tensor, $\bar{\bar{\sigma}} = \bar{\bar{\sigma}}^e + \bar{\bar{\sigma}}^m$, where $\bar{\bar{\sigma}}^e$ and $\bar{\bar{\sigma}}^m$ are the electric and magnetic field contributions respectively. $\bar{\bar{\sigma}}^e$ and $\bar{\bar{\sigma}}^m$ are given by:

$$\bar{\bar{\sigma}}^e(\tilde{\mathbf{r}}) = \varepsilon_o \left[\langle \mathbf{E}(\tilde{\mathbf{r}}, t) \mathbf{E}(\tilde{\mathbf{r}}, t) \rangle - \frac{1}{2} \bar{\bar{I}} \langle \mathbf{E}^2(\tilde{\mathbf{r}}, t) \rangle \right] \quad (8)$$

$$\bar{\bar{\sigma}}^m(\tilde{\mathbf{r}}) = \mu_o \left[\langle \mathbf{H}(\tilde{\mathbf{r}}, t) \mathbf{H}(\tilde{\mathbf{r}}, t) \rangle - \frac{1}{2} \bar{\bar{I}} \langle \mathbf{H}^2(\tilde{\mathbf{r}}, t) \rangle \right], \quad (9)$$

where $\langle \mathbf{E}(\tilde{\mathbf{r}}, t) \mathbf{E}(\tilde{\mathbf{r}}, t) \rangle$ and $\langle \mathbf{H}(\tilde{\mathbf{r}}, t) \mathbf{H}(\tilde{\mathbf{r}}, t) \rangle$ are matrices whose components are $\langle E_p(\tilde{\mathbf{r}}, t) E_q(\tilde{\mathbf{r}}, t) \rangle$ and $\langle H_p(\tilde{\mathbf{r}}, t) H_q(\tilde{\mathbf{r}}, t) \rangle$ respectively, $p, q = 1, 2, 3$, and $\bar{\bar{I}}$ is the identity matrix.

Since the fields are assumed to be stationary, $\langle \mathbf{E}(\tilde{\mathbf{r}}, t) \mathbf{E}(\tilde{\mathbf{r}}, t) \rangle$, $\langle \mathbf{H}(\tilde{\mathbf{r}}, t) \mathbf{H}(\tilde{\mathbf{r}}, t) \rangle$, and $\langle \mathbf{E}(\tilde{\mathbf{r}}, t) \mathbf{H}(\tilde{\mathbf{r}}, t) \rangle$ are independent of time [32]. The equal time correlation functions are related to the cross-spectral densities by:

$$\begin{aligned} \langle E_p(\tilde{\mathbf{r}}, t) E_q(\tilde{\mathbf{r}}, t) \rangle &= \int_0^\infty \frac{d\omega}{2\pi} \langle E_p(\tilde{\mathbf{r}}) E_q^*(\tilde{\mathbf{r}}) \rangle_s \\ &= \int_0^\infty \frac{d\omega}{2\pi} \langle E_p(\tilde{\mathbf{r}}) E_q^*(\tilde{\mathbf{r}}) + E_p^*(\tilde{\mathbf{r}}) E_q(\tilde{\mathbf{r}}) \rangle \end{aligned} \quad (10)$$

$$\begin{aligned} \langle H_p(\tilde{\mathbf{r}}, t) H_q(\tilde{\mathbf{r}}, t) \rangle &= \int_0^\infty \frac{d\omega}{2\pi} \langle H_p(\tilde{\mathbf{r}}) H_q^*(\tilde{\mathbf{r}}) \rangle_s \\ &= \int_0^\infty \frac{d\omega}{2\pi} \langle H_p(\tilde{\mathbf{r}}) H_q^*(\tilde{\mathbf{r}}) + H_p^*(\tilde{\mathbf{r}}) H_q(\tilde{\mathbf{r}}) \rangle \end{aligned} \quad (11)$$

$$\begin{aligned} \langle E_p(\tilde{\mathbf{r}}, t) H_q(\tilde{\mathbf{r}}, t) \rangle &= \int_0^\infty \frac{d\omega}{2\pi} \langle E_p(\tilde{\mathbf{r}}) H_q^*(\tilde{\mathbf{r}}) \rangle_s \\ &= \int_0^\infty \frac{d\omega}{2\pi} \langle E_p(\tilde{\mathbf{r}}) H_q^*(\tilde{\mathbf{r}}) + E_p^*(\tilde{\mathbf{r}}) H_q(\tilde{\mathbf{r}}) \rangle \end{aligned} \quad (12)$$

where the subscript s in Eq. 10 - Eq. 12 implies a symmetric sum. Using Eq. 1 and Eq. 2, we can express the cross-spectral densities of the components of the electric and magnetic field at $\tilde{\mathbf{r}} \in V_h$ as:

$$\langle \mathbf{E}(\tilde{\mathbf{r}}) \mathbf{E}^*(\tilde{\mathbf{r}}) \rangle_s = 2\omega\mu_o \left[\sum_{l=1}^N \Theta_l \bar{\bar{\mathcal{E}}}^{(l)}(\tilde{\mathbf{r}}) + \Theta_h \bar{\bar{\mathcal{E}}}^{(h)}(\tilde{\mathbf{r}}) \right] \quad (13)$$

$$\langle \mathbf{H}(\tilde{\mathbf{r}}) \mathbf{H}^*(\tilde{\mathbf{r}}) \rangle_s = 2\omega\varepsilon_o \left[\sum_{l=1}^N \Theta_l \bar{\bar{\mathcal{H}}}^{(l)}(\tilde{\mathbf{r}}) + \Theta_h \bar{\bar{\mathcal{H}}}^{(h)}(\tilde{\mathbf{r}}) \right] \quad (14)$$

$$\langle \mathbf{E}(\tilde{\mathbf{r}}) \mathbf{H}^*(\tilde{\mathbf{r}}) \rangle = \sum_{l=1}^N \Theta_l \bar{\bar{\mathcal{X}}}^{(l)}(\tilde{\mathbf{r}}) + \Theta_h \bar{\bar{\mathcal{X}}}^{(h)}(\tilde{\mathbf{r}}) \quad (15)$$

where $\Theta_l = \Theta(\omega, T_l)$, and $\langle \mathbf{E}(\tilde{\mathbf{r}}) \mathbf{E}^*(\tilde{\mathbf{r}}) \rangle_s$, $\langle \mathbf{H}(\tilde{\mathbf{r}}) \mathbf{H}^*(\tilde{\mathbf{r}}) \rangle_s$, and $\langle \mathbf{E}(\tilde{\mathbf{r}}) \mathbf{H}^*(\tilde{\mathbf{r}}) \rangle$ are matrices whose components are $\langle E_p(\tilde{\mathbf{r}}) E_q^*(\tilde{\mathbf{r}}) \rangle_s$, $\langle H_p(\tilde{\mathbf{r}}) H_q^*(\tilde{\mathbf{r}}) \rangle_s$, and $\langle E_p(\tilde{\mathbf{r}}) H_q^*(\tilde{\mathbf{r}}) \rangle$ respectively. Even though components of $\langle \mathbf{E} \mathbf{H}^* \rangle_s$ are necessary to compute radiative transfer, we persist with $\langle \mathbf{E} \mathbf{H}^* \rangle$. The reason for computing $\langle \mathbf{E} \mathbf{H}^* \rangle$ as opposed to $\langle \mathbf{E} \mathbf{H}^* \rangle_s$ will be clarified in Sec. 3.

The matrices $\bar{\bar{\mathcal{E}}}^{(l)}(\tilde{\mathbf{r}})$, $\bar{\bar{\mathcal{H}}}^{(l)}(\tilde{\mathbf{r}})$, and $\bar{\bar{\mathcal{X}}}^{(l)}(\tilde{\mathbf{r}})$ are contributions to $\langle \mathbf{E}(\tilde{\mathbf{r}}) \mathbf{E}^*(\tilde{\mathbf{r}}) \rangle_s$, $\langle \mathbf{H}(\tilde{\mathbf{r}}) \mathbf{H}^*(\tilde{\mathbf{r}}) \rangle_s$, and $\langle \mathbf{E}(\tilde{\mathbf{r}}) \mathbf{H}^*(\tilde{\mathbf{r}}) \rangle$ ($l = 1, 2, \dots, N, h$) from sources in volume V_l . For $l \in \{1, 2, \dots, N\}$, $\bar{\bar{\mathcal{E}}}^{(l)}(\tilde{\mathbf{r}})$, $\bar{\bar{\mathcal{H}}}^{(l)}(\tilde{\mathbf{r}})$, and $\bar{\bar{\mathcal{X}}}^{(l)}(\tilde{\mathbf{r}})$ are given by:

$$\begin{aligned} \bar{\bar{\mathcal{E}}}^{(l)}(\tilde{\mathbf{r}}) &= 2\Re \int_{V_l} d\mathbf{r} \left[\mu''(\mathbf{r}) \bar{\bar{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \bar{\bar{G}}_E^*(\mathbf{r}, \tilde{\mathbf{r}}) + \right. \\ &\quad \left. \varepsilon''(\mathbf{r}) |\mu(\mathbf{r})|^2 \frac{\omega^2}{c^2} \bar{\bar{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \bar{\bar{G}}_e^*(\mathbf{r}, \tilde{\mathbf{r}}) \right] \end{aligned} \quad (16a)$$

$$\begin{aligned} \bar{\bar{\mathcal{H}}}^{(l)}(\tilde{\mathbf{r}}) &= 2\Re \int_{V_l} d\mathbf{r} \left[\varepsilon''(\mathbf{r}) \bar{\bar{G}}_M^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \bar{\bar{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}}) + \right. \\ &\quad \left. \mu''(\mathbf{r}) |\varepsilon(\mathbf{r})|^2 \frac{\omega^2}{c^2} \bar{\bar{G}}_m^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \bar{\bar{G}}_m^*(\mathbf{r}, \tilde{\mathbf{r}}) \right] \end{aligned} \quad (16b)$$

$$\begin{aligned} \bar{\bar{\mathcal{X}}}^{(l)}(\tilde{\mathbf{r}}) &= i2 \frac{\omega^2}{c^2} \int_{V_l} d\mathbf{r} \left[\varepsilon''(\mathbf{r}) \mu(\mathbf{r}) \bar{\bar{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \bar{\bar{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}}) + \right. \\ &\quad \left. \mu''(\mathbf{r}) \varepsilon^*(\mathbf{r}) \bar{\bar{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \bar{\bar{G}}_m^*(\mathbf{r}, \tilde{\mathbf{r}}) \right] d\mathbf{r} \end{aligned} \quad (16c)$$

The expressions for $\overline{\mathcal{E}}^{(h)}(\tilde{\mathbf{r}})$, $\overline{\mathcal{H}}^{(h)}(\tilde{\mathbf{r}})$, and $\overline{\mathcal{X}}^{(h)}(\tilde{\mathbf{r}})$ have to be modified to take into account the singularity of the DGFs in the integrals in Eq. 17a-17c as $|\mathbf{r} - \tilde{\mathbf{r}}| \rightarrow 0$. The modified expressions for $\overline{\mathcal{E}}^{(h)}(\tilde{\mathbf{r}})$, $\overline{\mathcal{H}}^{(h)}(\tilde{\mathbf{r}})$, and $\overline{\mathcal{X}}^{(h)}(\tilde{\mathbf{r}})$ are:

$$\begin{aligned} \overline{\mathcal{E}}^{(h)}(\tilde{\mathbf{r}}) = & \lim_{V_\delta \rightarrow 0} 2\Re \int_{V_h} \left[\mu''(\mathbf{r}) \overline{\mathcal{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{G}}_E^*(\mathbf{r}, \tilde{\mathbf{r}}) \right. \\ & + \varepsilon''(\mathbf{r}) |\mu(\mathbf{r})|^2 \frac{\omega^2}{c^2} \overline{\mathcal{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{G}}_e^*(\mathbf{r}, \tilde{\mathbf{r}}) \Big] d\mathbf{r} \\ & + \varepsilon_h'' 2\Re \left(\frac{\mu_h}{\varepsilon_h^*} \overline{\mathcal{L}} \cdot \overline{\mathcal{G}}_e^{(sc)}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) + \frac{\mu_h}{\varepsilon_h^*} \overline{\mathcal{G}}_e^{(sc)T}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{L}} \right) \end{aligned} \quad (17a)$$

$$\begin{aligned} \overline{\mathcal{H}}^{(h)}(\tilde{\mathbf{r}}) = & \lim_{V_\delta \rightarrow 0} 2\Re \int_{V_h} \left[\varepsilon''(\mathbf{r}) \overline{\mathcal{G}}_M^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}}) \right. \\ & + \mu''(\mathbf{r}) |\varepsilon(\mathbf{r})|^2 \frac{\omega^2}{c^2} \overline{\mathcal{G}}_m^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{G}}_m^*(\mathbf{r}, \tilde{\mathbf{r}}) \Big] d\mathbf{r} \\ & + \mu_h'' 2\Re \left(\frac{\varepsilon_h}{\mu_h^*} \overline{\mathcal{L}} \cdot \overline{\mathcal{G}}_m^{(sc)}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) + \frac{\varepsilon_h}{\mu_h^*} \overline{\mathcal{G}}_m^{(sc)T}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{L}} \right) \end{aligned} \quad (17b)$$

$$\begin{aligned} \overline{\mathcal{X}}^{(h)}(\tilde{\mathbf{r}}) = & i2 \frac{\omega^2}{c^2} \lim_{V_\delta \rightarrow 0} \int_{V_i} \left[\varepsilon''(\mathbf{r}) \mu(\mathbf{r}) \overline{\mathcal{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}}) \right. \\ & + \mu''(\mathbf{r}) \varepsilon^*(\mathbf{r}) \overline{\mathcal{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{G}}_m^*(\mathbf{r}, \tilde{\mathbf{r}}) \Big] d\mathbf{r} \\ & - i2 \left(\frac{\varepsilon_h''}{\varepsilon_h} \overline{\mathcal{L}} \cdot \overline{\mathcal{G}}_M^{(sc)*}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) + \frac{\mu_h''}{\mu_h^*} \overline{\mathcal{G}}_E^{(sc)T}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) \cdot \overline{\mathcal{L}} \right) \end{aligned} \quad (17c)$$

where V_δ is volume of infinitesimal radius surrounding $\tilde{\mathbf{r}}$, $\overline{\mathcal{L}}$ is a shape dependent dyad [33], and $\overline{\mathcal{G}}_e(\mathbf{r}, \tilde{\mathbf{r}}) = \overline{\mathcal{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) + \overline{\mathcal{G}}_e^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$, $\overline{\mathcal{G}}_m(\mathbf{r}, \tilde{\mathbf{r}}) = \overline{\mathcal{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) + \overline{\mathcal{G}}_m^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$, $\overline{\mathcal{G}}_E(\mathbf{r}, \tilde{\mathbf{r}}) = \nabla \times \overline{\mathcal{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) + \overline{\mathcal{G}}_E^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$, and $\overline{\mathcal{G}}_M(\mathbf{r}, \tilde{\mathbf{r}}) = \nabla \times \overline{\mathcal{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) + \overline{\mathcal{G}}_M^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$. $\overline{\mathcal{G}}_o(\mathbf{r}, \tilde{\mathbf{r}})$ is the DGF when no scatterers are present, and $\overline{\mathcal{G}}^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$ is the contribution from presence of scatterers. $\overline{\mathcal{G}}^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$ is always finite. The volume integrals in Eq. 17a-Eq. 17c are finite even though $\varepsilon_h = \mu_h = 0$ because of the singularity in the DGFs. The last line of Eq. 17a-Eq. 17c are identically equal to zero for non-absorbing materials, including vacuum. However, we choose to retain them since they are essential for calculations of cross-spectral densities in absorbing media. We have neglected terms independent of the configuration of the scatterers. These terms are infinite because of the assumption that the thermal sources at any two locations are uncorrelated. They can be made finite by eliminating the local assumption in Eq. 1 but that will not affect the calculations of forces or heat transfer except at gaps smaller than the correlation length. Usually, the correlation length is of the order of the atomic spacing in dielectrics or the electron mean free path in metals. More detailed discussion of the singularity in DGFs and calculation of cross-spectral densities is given in Ref. [28].

3. Surface integral dyadic Green's function formalism

While the volume integrals in Eq. 16a - Eq. 17c can in principle be used to compute forces and radiative heat transfer, they are undesirable for the following reasons: (1) Evaluating classical radiative transfer between two objects requires the computation of the view factor between them. But for objects with simple geometries, computation of the view factor between two objects requires, in general, not the evaluation of a volume integral but the evaluation of a double integral over the surfaces of the two objects, (2) the expressions in Eq. 16a - Eq. 17c do not reflect the different reciprocity relations and boundary conditions satisfied by the DGFs, and (3) evaluation of volume integrals are computationally more expensive than that of surface integrals. These undesirable features can be overcome by converting Eq. 16a - Eq. 17c into appropriate surface integrals using Green's theorems for dyadic functions [34]. The surface integral representations for the cross-spectral densities are as follows:

$$\overline{\mathcal{E}}^{(l)}(\tilde{\mathbf{r}}) = 2\Im \oint_{S_l} [\mu(\mathbf{r}) \overline{\mathcal{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}})]^* \cdot [\mathbf{n}_l(\mathbf{r}) \times \overline{\mathcal{G}}_E(\mathbf{r}, \tilde{\mathbf{r}})] d\mathbf{r} \quad (18a)$$

$$\overline{\mathcal{H}}^{(l)}(\tilde{\mathbf{r}}) = 2\Im \oint_{S_l} [\varepsilon(\mathbf{r}) \overline{\mathcal{G}}_m^T(\mathbf{r}, \tilde{\mathbf{r}})]^* \cdot [\mathbf{n}_l(\mathbf{r}) \times \overline{\mathcal{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})] d\mathbf{r} \quad (18b)$$

$$\overline{\mathcal{X}}^{(l)}(\tilde{\mathbf{r}}) = - \oint_{S_l} [\overline{\mathcal{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot (\mathbf{n}_l(\mathbf{r}) \times \overline{\mathcal{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}})) + \quad (18c)$$

$$\frac{\omega^2}{c^2} \mu(\mathbf{r}) \overline{\mathcal{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot (\mathbf{n}_l(\mathbf{r}) \times \varepsilon(\mathbf{r}) \overline{\mathcal{G}}_m(\mathbf{r}, \tilde{\mathbf{r}}))^*] d\mathbf{r}$$

$$\begin{aligned} \overline{\mathcal{E}}^{(h)}(\tilde{\mathbf{r}}) = & -2 \sum_{l=1}^N \Im \oint_{S_l} \mu^*(\mathbf{r}) \overline{\mathcal{G}}_e^{*T}(\mathbf{r}, \tilde{\mathbf{r}}) \cdot (\mathbf{n}_l(\mathbf{r}) \times \overline{\mathcal{G}}_E(\mathbf{r}, \tilde{\mathbf{r}})) d\mathbf{r} \\ & + \Im(2\mu_h \overline{\mathcal{G}}_e(\tilde{\mathbf{r}}, \tilde{\mathbf{r}})) \end{aligned} \quad (18d)$$

$$\begin{aligned} \overline{\mathcal{H}}^{(h)}(\tilde{\mathbf{r}}) = & -2 \sum_{l=1}^N \Im \oint_{S_l} \varepsilon^*(\mathbf{r}) \overline{\mathcal{G}}_m^{*T}(\mathbf{r}, \tilde{\mathbf{r}}) \cdot (\mathbf{n}_l(\mathbf{r}) \times \overline{\mathcal{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})) d\mathbf{r} \\ & + \Im(2\varepsilon_h \overline{\mathcal{G}}_m(\tilde{\mathbf{r}}, \tilde{\mathbf{r}})) \end{aligned} \quad (18e)$$

$$\begin{aligned} \overline{\mathcal{X}}^{(h)}(\tilde{\mathbf{r}}) = & \sum_{l=1}^N \oint_{S_l} [\overline{\mathcal{G}}_E^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot (\mathbf{n}_l(\mathbf{r}) \times \overline{\mathcal{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}})) + \\ & \frac{\omega^2}{c^2} \mu(\mathbf{r}) \overline{\mathcal{G}}_e^T(\mathbf{r}, \tilde{\mathbf{r}}) \cdot (\mathbf{n}_l(\mathbf{r}) \times \varepsilon(\mathbf{r}) \overline{\mathcal{G}}_m(\mathbf{r}, \tilde{\mathbf{r}}))^*] d\mathbf{r} \\ & - i2\Im(\overline{\mathcal{G}}_M(\tilde{\mathbf{r}}, \tilde{\mathbf{r}})) \end{aligned} \quad (18f)$$

If the host medium is dissipative, the functions $\Im(2\mu_h \overline{\mathcal{G}}_e(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$, $\Im(2\varepsilon_h \overline{\mathcal{G}}_m(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$, and $\Im(\overline{\mathcal{G}}_M(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$ in Eq.

18d, Eq. 18e, and Eq. 18f respectively should be replaced by $\Im(2\mu_h \bar{G}_e^{(sc)}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$, $\Im(2\varepsilon_h \bar{G}_m^{(sc)}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$, and $\Im(\bar{G}_M^{(sc)}(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$. Using Eq. 18a-18f, Eq. 13-15 can be re-written as:

$$\langle \mathbf{E}\mathbf{E}^* \rangle_s = 2\omega\mu_o \sum_{l=1}^N (\Theta_l - \Theta_h) \bar{\mathcal{E}}^{(l)}(\tilde{\mathbf{r}}) + 2\omega\mu_o \Theta_h \Im(2\mu_h \bar{G}_e(\tilde{\mathbf{r}}, \tilde{\mathbf{r}})) \quad (19)$$

$$\langle \mathbf{H}\mathbf{H}^* \rangle_s = 2\omega\varepsilon_o \sum_{l=1}^N (\Theta_l - \Theta_h) \bar{\mathcal{H}}^{(l)}(\tilde{\mathbf{r}}) + 2\omega\varepsilon_o \Theta_h \Im(2\varepsilon_h \bar{G}_m(\tilde{\mathbf{r}}, \tilde{\mathbf{r}})) \quad (20)$$

$$\langle \mathbf{E}\mathbf{H}^* \rangle_s = \sum_{l=1}^N (\Theta_l - \Theta_h) \bar{\mathcal{X}}^{(l)}(\tilde{\mathbf{r}}) - i2\Theta_h \Im \bar{G}_M(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}) \quad (21)$$

The terms $\Im(2\mu_h \bar{G}_e(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$ and $\Im(2\varepsilon_h \bar{G}_m(\tilde{\mathbf{r}}, \tilde{\mathbf{r}}))$ are thermal equilibrium contributions to $\bar{\mathcal{E}}^{(h)}(\tilde{\mathbf{r}})$ and $\bar{\mathcal{H}}^{(h)}(\tilde{\mathbf{r}})$ respectively. They give rise to the van der Waals stresses as predicted by Lifshitz theory when $\varepsilon_h = \mu_h = 1$. The reason for persisting with $\langle \mathbf{E}\mathbf{H}^* \rangle_s$, as opposed to $\langle \mathbf{E}\mathbf{H}^* \rangle = \mathbf{E}\mathbf{H}^* + \mathbf{E}^*\mathbf{H}$, is to show that there is indeed an equilibrium contribution to $\langle \mathbf{E}\mathbf{H}^* \rangle_s$. However, when we compute $\langle \mathbf{E}\mathbf{H}^* \rangle_s$, the equilibrium contribution vanishes since radiative energy transfer between two objects at the same temperature must be zero.

Radiative transfer between two objects is discussed further in Sec. 4. Before proceeding to Sec. 4, we wish to remark on the form of the non-equilibrium contributions in Eq. 18a-18f. Using the property that $\bar{\mathbf{A}} = \mathbf{n}(\mathbf{n} \cdot \bar{\mathbf{A}}) - \mathbf{n} \times \mathbf{n} \times \bar{\mathbf{A}}$, we see that all the surface integrals in Eq. 18a-18f feature only tangential components of the dyadic Green's functions that are continuous across an interface between two materials.

4. Generalized transmissivity for radiative energy transfer

The steady state radiative heat transfer from object 1 to object 2 in Fig. 1, $Q_{1 \rightarrow 2}$, is given by:

$$Q_{1 \rightarrow 2} = - \oint_{S_2} \mathbf{P}^{(1)}(\tilde{\mathbf{r}}) \cdot \mathbf{n}_2(\tilde{\mathbf{r}}) d\tilde{\mathbf{r}} \quad (22)$$

where $\mathbf{P}^{(1)}(\tilde{\mathbf{r}})$ is the Poynting vector at $\tilde{\mathbf{r}} \in S_2$ due to thermally fluctuating sources within V_1 . The “-” sign in front of the surface integral is because $\mathbf{n}_2(\tilde{\mathbf{r}})$ is the outward pointing normal on the surface S_2 . The net heat transfer between objects 1 and 2 is given by $Q_{1,2} = Q_{1 \rightarrow 2} - Q_{2 \rightarrow 1}$. The components of $\mathbf{P}^{(1)}(\tilde{\mathbf{r}})$ are given by:

$$P_i^{(1)}(\tilde{\mathbf{r}}) = \epsilon_{iqp} \langle E_q(\tilde{\mathbf{r}}, t) H_p(\tilde{\mathbf{r}}, t) \rangle^{(1)}, \quad (23)$$

where $\langle E_q(\tilde{\mathbf{r}}, t) H_p(\tilde{\mathbf{r}}, t) \rangle^{(1)}$ is the contribution to $\langle E_q(\tilde{\mathbf{r}}, t) H_p(\tilde{\mathbf{r}}, t) \rangle$ from sources within V_1 . The

object indices 1 and 2 can be replaced by any $m, n \in \{1, 2, \dots, N\}$. From Sec. 2, we know that

$$\langle E_q(\tilde{\mathbf{r}}, t) H_p(\tilde{\mathbf{r}}, t) \rangle^{(1)} = \int_0^\infty \frac{d\omega}{2\pi} \langle E_q(\tilde{\mathbf{r}}, \omega) H_p^*(\tilde{\mathbf{r}}, \omega) \rangle_s^{(1)}.$$

Using Eq. 15, Eq. 21, Eq. 18c, and Eq. 23, Eq. 22 for $Q_{1 \rightarrow 2}$ can be re-written as:

$$\begin{aligned} Q_{1 \rightarrow 2} &= - \int_0^\infty \frac{d\omega}{2\pi} \Theta(\omega, T_1) \oint_{S_2} d\tilde{\mathbf{r}} n_{2i}(\tilde{\mathbf{r}}) \epsilon_{ipq} 2\Re \mathcal{X}_{pq}^{(1)}(\tilde{\mathbf{r}}) \\ &= \int_0^\infty \frac{d\omega}{2\pi} \Theta(\omega, T_1) T_{1 \rightarrow 2}^e(\omega) \\ &\Rightarrow T_{1 \rightarrow 2}^e(\omega) = - \oint_{S_2} d\tilde{\mathbf{r}} n_{2i}(\tilde{\mathbf{r}}) \epsilon_{ipq} 2\Re \mathcal{X}_{pq}^{(1)}(\tilde{\mathbf{r}}) \\ &= - \oint_{S_2} d\tilde{\mathbf{r}} \epsilon_{qip} n_{2i}(\tilde{\mathbf{r}}) 2\Re \mathcal{X}_{pq}^{(1)}(\tilde{\mathbf{r}}) \\ &= -2\Re \oint_{S_2} d\tilde{\mathbf{r}} \left(\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{\mathcal{X}}^{(1)}(\tilde{\mathbf{r}}) \right)_{qq} \\ &= -2\Re Tr \oint_{S_2} d\tilde{\mathbf{r}} \left(\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{\mathcal{X}}^{(1)}(\tilde{\mathbf{r}}) \right), \end{aligned} \quad (24)$$

where $T_{1 \rightarrow 2}^e(\omega)$ is a generalized transmissivity for radiative energy transport between objects 1 and 2, and $Tr(\bar{\mathbf{A}}) = \sum_{p=1}^3 A_{pp}$. The superscript e in $T_{1 \rightarrow 2}^e(\omega)$ stands for “energy.”

Substituting the expression for $\bar{\mathcal{X}}^{(1)}$ from Eq. 18c in the last line of Eq. 24, $T_{1 \rightarrow 2}^e(\omega)$ can be shown to be:

$$\begin{aligned} T_{1 \rightarrow 2}^e(\omega) &= 2\Re Tr \oint_{S_1} d\mathbf{r} \oint_{S_2} d\tilde{\mathbf{r}} \left[\frac{\omega^2}{c^2} \times \right. \\ &\quad \left. [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \mu_2 \bar{G}_e(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \varepsilon_1^* \bar{G}_m^*(\mathbf{r}, \tilde{\mathbf{r}})] \right. \\ &\quad \left. + [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{G}_E(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \bar{G}_E^*(\mathbf{r}, \tilde{\mathbf{r}})] \right] \end{aligned} \quad (25)$$

where $\hat{\mathbf{n}}_1(\mathbf{r})$ is the outward pointing normal on the surface S_1 , as shown in Fig. 1. For any two vectors $\mathbf{a}, \mathbf{b}$ and dyads $\bar{\mathbf{A}}, \bar{\mathbf{B}}$, the following property can be shown to be true: $Tr\{(\mathbf{a} \times \bar{\mathbf{A}}) \cdot (\mathbf{b} \times \bar{\mathbf{B}})\} = Tr\{(\mathbf{b} \times \bar{\mathbf{A}}^T) \cdot (\mathbf{a} \times \bar{\mathbf{B}}^T)\}$. Using this property, and the reciprocity relations (Eq. 4, Eq. 5, and Eq. 6), we can derive the following equations:

$$\begin{aligned} \Re Tr[\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \mu_2 \bar{G}_e(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \varepsilon_1^* \bar{G}_m^*(\mathbf{r}, \tilde{\mathbf{r}})] &= \\ \Re Tr[\hat{\mathbf{n}}_1(\mathbf{r}) \times \mu_1 \bar{G}_e(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \varepsilon_2^* \bar{G}_m^*(\tilde{\mathbf{r}}, \mathbf{r})] \end{aligned} \quad (26a)$$

$$\begin{aligned} \Re Tr[\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{G}_E(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \bar{G}_E^*(\mathbf{r}, \tilde{\mathbf{r}})] &= \\ \Re Tr[\hat{\mathbf{n}}_1(\mathbf{r}) \times \bar{G}_M(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{G}_M^*(\tilde{\mathbf{r}}, \mathbf{r})] \end{aligned} \quad (26b)$$

With the aid of Eq. 26, $T_{1 \rightarrow 2}^e(\omega)$ can also be shown to be:

$$T_{1 \rightarrow 2}^e(\omega) = 2\Re Tr \oint_{S_1} d\mathbf{r} \oint_{S_2} d\tilde{\mathbf{r}} \left[\frac{\omega^2}{c^2} \times \right. \\ \left. [\hat{\mathbf{n}}_1(\mathbf{r}) \times \mu_1 \overline{\overline{G}}_e(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \varepsilon_2^* \overline{\overline{G}}_m^*(\tilde{\mathbf{r}}, \mathbf{r})] \right. \\ \left. + [\hat{\mathbf{n}}_1(\mathbf{r}) \times \overline{\overline{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \overline{\overline{G}}_M^*(\tilde{\mathbf{r}}, \mathbf{r})] \right] \quad (27)$$

The generalized transmissivity from object 2 to object 1, $T_{2 \rightarrow 1}^e(\omega)$ can be determined from Eq. 27 (or Eq. 25) by interchanging the subscripts 1 and 2 ($\mathbf{r} \in S_1$ and $\tilde{\mathbf{r}} \in S_2$ are dummy variables and do not affect the value of the double integral). $T_{2 \rightarrow 1}^e(\omega)$ is given by:

$$T_{2 \rightarrow 1}^e(\omega) = 2\Re Tr \oint_{S_2} d\tilde{\mathbf{r}} \oint_{S_1} d\mathbf{r} \left[\frac{\omega^2}{c^2} \times \right. \\ \left. [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \mu_2 \overline{\overline{G}}_e(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \varepsilon_1^* \overline{\overline{G}}_m^*(\mathbf{r}, \tilde{\mathbf{r}})] \right. \\ \left. + [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \overline{\overline{G}}_M(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \overline{\overline{G}}_M^*(\mathbf{r}, \tilde{\mathbf{r}})] \right] \quad (28)$$

That the expressions for generalized transmissivity derived earlier (Eq. 25 or Eq. 27) satisfy the principle of reciprocity in thermal radiative transfer, i.e. $T_{2 \rightarrow 1}^e(\omega) = T_{1 \rightarrow 2}^e(\omega)$, can be established by using Eq. 26b to modify the expression for $T_{2 \rightarrow 1}^e(\omega)$ as follows:

$$T_{2 \rightarrow 1}^e(\omega) = 2\Re Tr \oint_{S_2} d\tilde{\mathbf{r}} \oint_{S_1} d\mathbf{r} \left[\frac{\omega^2}{c^2} \times \right. \\ \left. [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \mu_2 \overline{\overline{G}}_e(\tilde{\mathbf{r}}, \mathbf{r})] \cdot [\hat{\mathbf{n}}_1(\mathbf{r}) \times \varepsilon_1^* \overline{\overline{G}}_m^*(\mathbf{r}, \tilde{\mathbf{r}})] \right. \\ \left. + [\hat{\mathbf{n}}_1(\mathbf{r}) \times \overline{\overline{G}}_E(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \overline{\overline{G}}_E^*(\tilde{\mathbf{r}}, \mathbf{r})] \right] \\ = T_{1 \rightarrow 2}^e(\omega) \quad (\text{Eq. 25})$$

Though expressions for $T_{1 \rightarrow 2}^e(\omega)$ in Eq. 25 and Eq. 27 are surface integrals, they are in fact derived from a volumetric integral over V_1 (Eq. 16c). Similarly, energy emission from the object V_l ($l = 1, 2, \dots, N$) is derived from a volumetric integration over V_l . For this reason, the formulae derived for $T_{1 \rightarrow 2}^e(\omega)$ and $T_{2 \rightarrow 1}^e(\omega)$ (Eq. 25, Eq. 27, or Eq. 29) can be described as “interior formulae.” The integration over V_1 (for $T_{1 \rightarrow 2}^e(\omega)$) or V_2 (for $T_{2 \rightarrow 1}^e(\omega)$) is made explicit by the presence of ε_1, μ_2 (in Eq. 25) or ε_2, μ_1 (Eq. 27, Eq. 29). The corresponding “exterior formula” should not involve, or appear not to involve, any of these properties in the formula for transmissivity. The exterior formula for $T_{1 \rightarrow 2}^e$ can be derived by using the boundary conditions (Eq. 3a-Eq. 3d) and converting Eq. 27 into the following equation:

$$T_{1 \rightarrow 2}^e(\omega) = 2\Re Tr \oint_{S_1} d\mathbf{r} \oint_{S_2} d\tilde{\mathbf{r}} \left[\frac{\omega^2}{c^2} \times \right. \\ \left. [\hat{\mathbf{n}}_1(\mathbf{r}) \times \mu_h \overline{\overline{G}}_e(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \varepsilon_h^* \overline{\overline{G}}_m^*(\tilde{\mathbf{r}}, \mathbf{r})] \right. \\ \left. + [\hat{\mathbf{n}}_1(\mathbf{r}) \times \overline{\overline{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})] \cdot [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \overline{\overline{G}}_M^*(\tilde{\mathbf{r}}, \mathbf{r})] \right] \quad (30)$$

There is a correspondence between the “direct” and “indirect” methods [35, 36, 37] and the “interior formula” and “exterior formula” derived above. It can be shown that the “exterior formula” is a generalization of the “indirect” method to include problems of near-field thermal radiative energy transfer between two objects, in addition to the calculation of thermal emission from objects for which it is currently used. This correspondence will be undertaken in a future work and is not pursued any further in this paper.

Since $T_{2 \rightarrow 1}^e(\omega) = T_{1 \rightarrow 2}^e(\omega)$, the net radiative exchange between objects 1 and 2, $Q_{1,2}$, is given by:

$$Q_{1,2} = \int_0^\infty \frac{d\omega}{2\pi} [\Theta(\omega, T_1) - \Theta(\omega, T_2)] T_{1 \rightarrow 2}^e(\omega) \quad (31)$$

From Eq. 31, a linearized conductance for radiative transfer between objects 1 and 2 can be defined as:

$$G_{1,2}^e(T) = \lim_{T_1, T_2 \rightarrow T} \frac{Q_{1,2}}{T_1 - T_2} = \int_0^\infty \frac{d\omega}{2\pi} \frac{\partial \Theta}{\partial T} T_{1 \rightarrow 2}^e(\omega) \quad (32)$$

4.1. Generalized transmissivity for radiative momentum transfer?

Though expressions for generalized transmissivity in terms of DGFs (Eq. 25, Eq. 27, Eq. 29, and Eq. 30) have been derived for energy transfer, we have been unable to obtain equivalent expressions for generalized (vectorial) transmissivity or conductance of thermal non-equilibrium momentum transfer. Why this is so can be explained by considering the nature of the Poynting vector and the electromagnetic stress tensor. It is a well-known property of Maxwell’s equations that the electric field at any location due to sources within a particular object, for instance object 1 in Fig. 1, can be expressed in terms of surface integrals of tangential electric and magnetic fields on the surface of that object. This property of electromagnetic fields forms the basis for the boundary element method for numerical solution of electromagnetic scattering problems. The normal component of the Poynting vector on the surface of object 2, $(\mathbf{E}(\tilde{\mathbf{r}}, t) \times \mathbf{H}(\tilde{\mathbf{r}}, t)) \cdot \mathbf{n}_2(\tilde{\mathbf{r}})$, has an additional property that it can be written as $[(\mathbf{n}_2(\tilde{\mathbf{r}}) \times \mathbf{E}(\tilde{\mathbf{r}}, t)) \times (\mathbf{n}_2(\tilde{\mathbf{r}}) \times \mathbf{H}(\tilde{\mathbf{r}}, t))] \cdot \mathbf{n}_2(\tilde{\mathbf{r}})$. This ensures that the *radiative heat transfer between objects 1 and 2 can be expressed in terms of tangential electric and magnetic fields on the surfaces of both objects*. This property of radiative heat transfer is reflected in the different expressions for $T_{1 \rightarrow 2}^e(\omega)$ because they contain only tangential components of the DGFs on the surfaces of both objects. However, the electric and magnetic stress tensors in Eq. 8 and Eq. 9 do not share this property. The force exerted by object 1 on object 2 requires knowledge of not only the tangential components of $\mathbf{E}$ and $\mathbf{H}$ on the surface of object 2 but also the normal components. While our inability to deduce an appropriate form for the transmissivity for momentum transfer does not mean that such a

transmissivity does not exist, Eq. 19, and Eq. 20 can still be used to determine $\overline{\sigma}^e(\mathbf{r})$ (Eq. 8) and $\overline{\sigma}^m(\mathbf{r})$ (Eq. 9) for specific geometric configuration of objects, from which thermal non-equilibrium van der Waals forces between objects can be computed.

5. Application to specific geometries or properties

5.1. Planar multilayered media

Since theoretical analysis of radiative transfer and thermal non-equilibrium van der Waals forces between planar multilayered objects have been published in literature, the expressions derived in Sec. 3 are applied to objects shown in Fig. 2 and the resultant expressions compared with those in literature. The half spaces in Fig. 2, marked 1 and 2, can be homogeneous materials or can be composed of planar multilayer films. The only requirement is that the temperature gradients within 1 and 2 are negligible enough that they can be approximated as thermal reservoirs at temperatures T_1 and T_2 . The vacuum layer of thickness l separating the two objects is equivalent to the host medium in Fig. 1. The planar media lie in the $x - y$ plane and the unit vector in the z direction, $\hat{z}$, is directed from object 1 to object 2. The interfaces of objects 1 and 2 with vacuum are at coordinates $z = z_1$ and $z = z_2$ ($|z_2 - z_1| = l$). The polarization dependent reflection coefficient of electromagnetic plane wave originating in vacuum and incident at the surface of half space 1 (in the absence of half space 2) is denoted as $\tilde{R}_{h1}^{(\mu)}$, where $\mu = s$ (transverse electric), p (transverse magnetic). A similar reflection coefficient for waves incident on half space 2, in the absence of half space 1, is denoted $\tilde{R}_{h2}^{(\mu)}$. $\overline{G}_o(\mathbf{r}, \tilde{\mathbf{r}})$ for any two locations $\mathbf{r}$ and $\tilde{\mathbf{r}}$ within the vacuum layer is given by:

$$\overline{G}_o(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{i}{4\pi} \int \frac{dk_\rho k_\rho}{k_{hz}} e^{i\mathbf{k}_\rho \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}')} \times \sum_{\mu=s,p} \begin{cases} \hat{x}^{(\mu)}(+k_{hz}) \hat{x}^{(\mu)}(+k_{hz}) e^{i(z-z')k_{hz}} & \text{if } z > z' \\ \hat{x}^{(\mu)}(-k_{hz}) \hat{x}^{(\mu)}(-k_{hz}) e^{i(z'-z)k_{hz}} & \text{if } z < z' \end{cases} \quad (33)$$

where $\mathbf{k}_\rho = k_\rho \hat{k}_\rho = k_x \hat{x} + k_y \hat{y}$, $\hat{x}^{(s)}(\pm k_{hz}) = \hat{k}_\rho \times \hat{z} = (k_y \hat{x} - k_x \hat{y})/k_\rho$, $\hat{x}^{(p)}(\pm k_{hz}) = (\mp k_{hz} \hat{k}_\rho + k_\rho \hat{z})/k_h$, and $k_{hz}^2 + k_\rho^2 = k_h^2$. For vacuum, $\mu_h = \varepsilon_h = 1$ and $k_h = \omega/c$. The scattered DGF, $\overline{G}_e^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$, is given by:

$$\overline{G}_e^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{i}{4\pi} \int \frac{dk_\rho k_\rho}{k_{hz}} \sum_{\mu=s,p} \frac{e^{i\mathbf{k}_\rho \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}')}}{D^{(\mu)}} \times \sum_{\nu=\pm 1} \sum_{\xi=\pm 1} C_{\nu,\xi}^{(\mu)} \hat{x}^{(\mu)}(\nu k_{hz}) \hat{x}^{(\mu)}(\xi k_{hz}) e^{i(\nu z - \xi z')k_{hz}}, \quad (34)$$

where

$$D^{(\mu)} = 1 - \tilde{R}_{h1}^{(\mu)} \tilde{R}_{h2}^{(\mu)} e^{i2k_{hz}l} \quad (35)$$

and

$$C_{\nu,\xi}^{(\mu)} = \begin{cases} \tilde{R}_{h1}^{(\mu)} \tilde{R}_{h2}^{(\mu)} e^{i2k_{hz}l} & \text{if } \nu = \xi \\ \tilde{R}_{h2}^{(\mu)} e^{i2k_{hz}z_2} & \text{if } \nu = -1, \xi = 1 \\ \tilde{R}_{h1}^{(\mu)} e^{-i2k_{hz}z_1} & \text{if } \nu = 1, \xi = -1 \end{cases} \quad (36)$$

The total DGF, $\overline{G}_e(\mathbf{r}, \tilde{\mathbf{r}})$, is given by $\overline{G}_o(\mathbf{r}, \tilde{\mathbf{r}}) + \overline{G}_e^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$. The reflection coefficients $\tilde{R}_{h1}^{(\mu)}$ and $\tilde{R}_{h2}^{(\mu)}$ at the interface between two homogeneous media are given by the usual Fresnel reflection coefficients. For multilayered media, they can be computed using the transfer matrix method or by using recursion relations [38]. Since both $\mathbf{r}$ and $\tilde{\mathbf{r}}$ are within the same layer, $\overline{G}_m(\mathbf{r}, \tilde{\mathbf{r}})$ and $\overline{G}_M(\mathbf{r}, \tilde{\mathbf{r}})$ can be determined by simply replacing all occurrences of $\varepsilon(\omega)$ by the corresponding $\mu(\omega)$ and vice versa. Explicit expressions for $\overline{G}_m^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$, $\overline{G}_E^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$, and $\overline{G}_M^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}})$ are given below:

$$\overline{G}_m^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{i}{4\pi} \int \frac{dk_\rho k_\rho}{k_{hz}} \sum_{\mu=s,p} \frac{e^{i\mathbf{k}_\rho \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}')}}{D^{(\mu')}} \times \sum_{\nu=\pm 1} \sum_{\xi=\pm 1} C_{\nu,\xi}^{(\mu')} \hat{x}^{(\mu)}(\nu k_{hz}) \hat{x}^{(\mu)}(\xi k_{hz}) e^{i(\nu z - \xi z')k_{hz}}, \quad (37)$$

where $\mu' = p$ if $\mu = s$ and $\mu' = s$ if $\mu = p$. Defining $\hat{k}_h^{(\pm)} = (k_\rho \hat{k}_\rho \pm k_{hz} \hat{z})/k_h$, we have the following relations: $\hat{k}_h^{(\pm)} \times \hat{x}^{(s)}(\pm k_{hz}) = -\hat{x}^{(p)}(\pm k_{hz})$ and $\hat{k}_h^{(\pm)} \times \hat{x}^{(p)}(\pm k_{hz}) = \hat{x}^{(s)}(\pm k_{hz})$. Using these relations, we obtain:

$$\overline{G}_E^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{-1}{4\pi} \int \frac{dk_\rho k_\rho}{k_{hz}} k_h \sum_{\mu=s,p} \frac{e^{i\mathbf{k}_\rho \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}')}}{D^{(\mu)}} \times \sum_{\nu=\pm 1} \sum_{\xi=\pm 1} (-1)^\beta C_{\nu,\xi}^{(\mu)} \hat{x}^{(\mu')}(\nu k_{hz}) \hat{x}^{(\mu)}(\xi k_{hz}) e^{i(\nu z - \xi z')k_{hz}}, \quad (38)$$

$$\overline{G}_M^{(sc)}(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{-1}{4\pi} \int \frac{dk_\rho k_\rho}{k_{hz}} k_h \sum_{\mu=s,p} \frac{e^{i\mathbf{k}_\rho \cdot (\boldsymbol{\rho} - \boldsymbol{\rho}')}}{D^{(\mu')}} \times \sum_{\nu=\pm 1} \sum_{\xi=\pm 1} (-1)^\beta C_{\nu,\xi}^{(\mu')} \hat{x}^{(\mu')}(\nu k_{hz}) \hat{x}^{(\mu)}(\xi k_{hz}) e^{i(\nu z - \xi z')k_{hz}}, \quad (39)$$

where $\beta = 0$ if $\mu = p$ and $\beta = 1$ if $\mu = s$.

A key distinction between the “direct” and “exterior” methods can be illustrated through the example being considered here. Let us assume that the half space L is composed of multiple planar films. In the direct method [35, 36], the contribution of each layer is evaluated separately and added subsequently to determine the heat flux in the vacuum layer due to half space L . This requires finding the DGFs when $\tilde{\mathbf{r}}$ belongs to the vacuum layer and $\mathbf{r}$ lies in each of the thin films that makes up the half spaces L . On the other hand, the exterior method, which is used to derive Eq. 40 and Eq. 41, requires only knowledge of the DGFs when $\mathbf{r}$ and $\tilde{\mathbf{r}}$ belong to the vacuum layer. Though




Figure 2: Two multilayered half spaces at T_1 and T_2 separated by a vacuum gap.

one can derive the same results, but with more algebraic manipulations, for layered media, the ease of using the exterior method should become apparent when one tries to model near-field radiative transfer in more complicated geometries, for instance between two coated spheres.

5.1.1. Two parallel half spaces: Radiative transfer

For the two half spaces in Fig. 2, the surface normal vectors $\hat{\mathbf{n}}(\mathbf{r})$ and $\hat{\mathbf{n}}(\tilde{\mathbf{r}})$ in Eq. 30 are given by $\hat{z}$ and $-\hat{z}$ respectively. $\bar{G}_e(\mathbf{r}, \tilde{\mathbf{r}})$, $\bar{G}_m(\mathbf{r}, \tilde{\mathbf{r}})$, $\bar{G}_E(\mathbf{r}, \tilde{\mathbf{r}})$, and $\bar{G}_M(\mathbf{r}, \tilde{\mathbf{r}})$ derived using Eq. 33 and Eq. 34 are substituted into Eq. 30 to obtain, $T_{1 \rightarrow 2}^{e,pp}$, the generalized transmissivity of energy transfer between planar half spaces (the superscript pp is short for “planar-planar” and is used to indicate the type of objects). After some manipulations, $T_{1 \rightarrow 2}^{e,pp}$ can be shown to be:

$$T_{1 \rightarrow 2}^{e,pp}(\omega) = \int_0^{\omega/c} \frac{k_\rho dk_\rho}{2\pi} \sum_{\mu=s,p} \frac{(1 - |\tilde{R}_{h1}^{(\mu)}|^2)(1 - |\tilde{R}_{h2}^{(\mu)}|^2)}{|1 - \tilde{R}_{h1}^{(\mu)} \tilde{R}_{h2}^{(\mu)} e^{i2k_{hz}l}|^2} + \int_{\omega/c}^{\infty} \frac{k_\rho dk_\rho}{2\pi} \sum_{\mu=s,p} \frac{4\Im(\tilde{R}_{h1}^{(\mu)})\Im(\tilde{R}_{h2}^{(\mu)})e^{-2|k_{hz}l|}}{|1 - \tilde{R}_{h1}^{(\mu)} \tilde{R}_{h2}^{(\mu)} e^{-2|k_{hz}l|}|^2} \quad (40)$$

This expression for $T_{1 \rightarrow 2}^{e,pp}(\omega)$ is in agreement with expressions for transmissivity of energy transfer across half spaces [9]. For the case of two homogeneous half spaces, we have also confirmed that volume integral expression (Eq. 16c) as well as surface integral expressions (Eq. 25, Eq. 30) yield the same result. Two interesting features of Eq. 40 need to be emphasized: (1) Eq. 40 is valid for energy transfer not just between two homogeneous half spaces but also between two half spaces comprising planar thin films, (2) It is valid for isotropic materials with electric as well as magnetic polarizabilities, i.e., with frequency dependent ϵ and μ .

5.1.2. Two parallel half spaces: Non-equilibrium pressure

Since we have not been able to derive a generalized transmissivity for momentum transfer, we use Eq. 8 and

Eq. 9 to derive the van der Waals pressure in the vacuum gap. Because the film is perpendicular to the z direction, the van der Waals pressure is given by the zz component of the stress tensor, σ_{zz} . Using the expressions for $\bar{G}_e(\mathbf{r}, \tilde{\mathbf{r}})$ and $\bar{G}_E(\mathbf{r}, \tilde{\mathbf{r}})$ in Eq. 19 and $\bar{G}_m(\mathbf{r}, \tilde{\mathbf{r}})$ and $\bar{G}_M(\mathbf{r}, \tilde{\mathbf{r}})$ in Eq. 20, a *generalized transmissivity for momentum flux* from L to R is given by:

$$T_{1 \rightarrow 2}^{m,pp}(\omega) = - \int_0^{\omega/c} \frac{k_\rho dk_\rho}{2\pi} \frac{k_{hz}}{\omega} \sum_{\mu=s,p} \frac{(1 - |\tilde{R}_{h1}^{(\mu)}|^2)(1 + |\tilde{R}_{h2}^{(\mu)}|^2)}{|1 - \tilde{R}_{h1}^{(\mu)} \tilde{R}_{h2}^{(\mu)} e^{i2k_{hz}l}|^2} + \int_{\omega/c}^{\infty} \frac{k_\rho dk_\rho}{2\pi} \frac{|k_{hz}|}{\omega} \sum_{\mu=s,p} \frac{4\Im(\tilde{R}_{h1}^{(\mu)})\Re(\tilde{R}_{h2}^{(\mu)})e^{-2|k_{hz}l|}}{|1 - \tilde{R}_{h1}^{(\mu)} \tilde{R}_{h2}^{(\mu)} e^{-2|k_{hz}l|}|^2} \quad (41)$$

The superscript m in $T_{1 \rightarrow 2}^{m,pp}(\omega)$ stands for “momentum.” Using the same notation as Antezza et. al. [39], the non-equilibrium pressure in the vacuum layer due to temperature T_L of half space L while $T_R = 0$ K, denoted by $P_{neq}(T_L, T_R = 0, l)$, is calculated using the formula $P_{neq}(T_L, T_R = 0, l) = \int_0^{\infty} \frac{d\omega}{2\pi} \Theta(\omega, T_L) T_{1 \rightarrow 2}^{m,pp}(\omega)$. It is interesting to note that Eq. 41, which is valid for half spaces with arbitrary ϵ and μ , coincide with the expressions for non-equilibrium van der Waals pressure and radiative transfer derived in Ref. [39] and Ref. [9] even though the authors of Ref. [39] and Ref. [9] derived it only for the case when $\mu = 1$ everywhere.

5.2. Agreement with theory of blackbody radiative transfer

Computing radiative transfer between two arbitrarily shaped isotropic objects using Eq. 25, Eq. 27, or Eq. 30 is computationally involved because of the need to compute the appropriate DGFs. However, for one class of objects, namely blackbodies (or objects those can be approximated as blackbodies), the expression for generalized transmissivity derived here can be used to obtain useful results irrespective of the shape. Blackbody radiative transfer is derived from Planck’s theory of blackbody radiation and Kirchoff’s laws, both of which are consequences of thermodynamics applied to relatively simple electrodynamical systems (for example, photon gas in a piston with perfectly reflective walls). Using thermodynamic arguments, this idea is generalized to arbitrarily shaped objects to yield $Q_{1,2}^{bb} = A_1 F_{1,2} \sigma_{SB} (T_1^4 - T_2^4)$, where σ_{SB} is the Stefan-Boltzmann constant, the superscript bb stands for “blackbody”, and $F_{1,2}$ is the view factor between objects 1 and 2. The view factor $F_{1,2}$ between the two objects in Fig. 1 is given by:

$$F_{1,2} = \frac{1}{A_1} \oint_{S_1} d\mathbf{r} \oint_{S_2} d\tilde{\mathbf{r}} \frac{(-\mathbf{n}_1 \cdot \hat{\mathbf{R}})(\mathbf{n}_2 \cdot \hat{\mathbf{R}})}{\pi R^2}, \quad (42)$$

where $\hat{\mathbf{R}} = (\mathbf{r} - \tilde{\mathbf{r}})/|\mathbf{r} - \tilde{\mathbf{r}}|$, $\mathbf{R} = R\hat{\mathbf{R}} = \mathbf{r} - \tilde{\mathbf{r}}$, $\mathbf{r} \in S_1$, and $\tilde{\mathbf{r}} \in S_2$, and A_1 is the area of S_1 .

A blackbody is one that absorbs all radiation incident on it and scatters none. For an object in vacuum ($\varepsilon_h = \mu_h = 1$), this can be achieved by a region of space (the blackbody) with permittivity and permeability given by $\varepsilon = 1 + i\delta$, $\mu = 1 + i\gamma$ such that $\delta, \gamma \rightarrow 0$, ensuring that there is no scattering by the object. The nominal dimension L should be such that $\delta kL \gg 1$ or $\gamma kL \gg 1$, where $k = \omega/c$ and c is the speed of light in vacuum, ensuring that all the radiation entering the object is absorbed. Because the properties of the objects differ infinitesimally from that of the host medium, scattering can effectively be neglected and the DGFs, $\bar{\bar{G}}_e(\mathbf{r}, \tilde{\mathbf{r}})$ and $\bar{\bar{G}}_m(\mathbf{r}, \tilde{\mathbf{r}})$, are simply given by the DGF in free space, which is:

$$\bar{\bar{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{e^{ik_h R}}{4\pi R} \left[\hat{\mathbf{R}}\hat{\mathbf{R}} \left(-i\frac{2}{k_h R} + \frac{2}{k_h^2 R^2} \right) + (\bar{\bar{I}} - \hat{\mathbf{R}}\hat{\mathbf{R}}) \left(1 + \frac{i}{k_h R} - \frac{1}{k_h^2 R^2} \right) \right] \quad (43)$$

When the spacing between objects is large compared to the wavelength, Eq. 43 reduces to:

$$\bar{\bar{G}}_e(\mathbf{r}, \tilde{\mathbf{r}}) = \bar{\bar{G}}_m(\mathbf{r}, \tilde{\mathbf{r}}) = \bar{\bar{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) = \frac{e^{ik_h R}}{4\pi R} (\bar{\bar{I}} - \hat{\mathbf{R}}\hat{\mathbf{R}}) \quad (44)$$

Similarly, $\bar{\bar{G}}_E(\mathbf{r}, \tilde{\mathbf{r}})$ and $\bar{\bar{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})$ can be written as:

$$\nabla \times \bar{\bar{G}}_o(\mathbf{r}, \tilde{\mathbf{r}}) = \bar{\bar{G}}_O(\mathbf{r}, \tilde{\mathbf{r}}) = ik_h \frac{e^{ik_h R}}{4\pi R} (\hat{\mathbf{R}} \times \bar{\bar{I}}) \quad (45)$$

To derive the generalized transmissivity between two blackbodies, the following derivations are useful:

$$\begin{aligned} \Re Tr[(\hat{\mathbf{n}}_1(\mathbf{r}) \times \mu_h \bar{\bar{G}}_e(\mathbf{r}, \tilde{\mathbf{r}})) \cdot (\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \varepsilon_h \bar{\bar{G}}_m(\tilde{\mathbf{r}}, \mathbf{r}))^*] \\ = \Re Tr[(\hat{\mathbf{n}}_1(\mathbf{r}) \times \bar{\bar{G}}_o(\mathbf{r}, \tilde{\mathbf{r}})) \cdot (\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{\bar{G}}_o(\tilde{\mathbf{r}}, \mathbf{r}))^*] \\ = \frac{[\hat{\mathbf{n}}_1(\mathbf{r}) \times (\bar{\bar{I}} - \hat{\mathbf{R}}\hat{\mathbf{R}})]_{ij} [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times (\bar{\bar{I}} - \hat{\mathbf{R}}\hat{\mathbf{R}})]_{ji}}{(4\pi R)^2} \\ = \frac{[\hat{\mathbf{n}}_1(\mathbf{r}) \times (\hat{\theta}\hat{\theta} + \hat{\phi}\hat{\phi})]_{ij} [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times (\hat{\theta}\hat{\theta} + \hat{\phi}\hat{\phi})]_{ji}}{(4\pi R)^2} \\ = \frac{-2(\hat{\mathbf{n}}_1(\mathbf{r}) \cdot \hat{\mathbf{R}})(\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \cdot \hat{\mathbf{R}})}{(4\pi R)^2} \end{aligned} \quad (46)$$

$$\begin{aligned} \Re Tr[(\hat{\mathbf{n}}_1(\mathbf{r}) \times \bar{\bar{G}}_M(\mathbf{r}, \tilde{\mathbf{r}})) \cdot (\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{\bar{G}}_M(\tilde{\mathbf{r}}, \mathbf{r}))^*] \\ = \Re Tr[(\hat{\mathbf{n}}_1(\mathbf{r}) \times \bar{\bar{G}}_O(\mathbf{r}, \tilde{\mathbf{r}})) \cdot (\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times \bar{\bar{G}}_O(\tilde{\mathbf{r}}, \mathbf{r}))^*] \\ = -k_h^2 \frac{[\hat{\mathbf{n}}_1(\mathbf{r}) \times (\hat{\mathbf{R}} \times \bar{\bar{I}})]_{ij} [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times (\hat{\mathbf{R}} \times \bar{\bar{I}})]_{ji}}{(4\pi R)^2} \\ = -k_h^2 \frac{[\hat{\mathbf{n}}_1(\mathbf{r}) \times (\hat{\phi}\hat{\theta} - \hat{\theta}\hat{\phi})]_{ij} [\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \times (\hat{\phi}\hat{\theta} - \hat{\theta}\hat{\phi})]_{ji}}{(4\pi R)^2} \\ = k_h^2 \frac{-2(\hat{\mathbf{n}}_1(\mathbf{r}) \cdot \hat{\mathbf{R}})(\hat{\mathbf{n}}_2(\tilde{\mathbf{r}}) \cdot \hat{\mathbf{R}})}{(4\pi R)^2} \end{aligned} \quad (47)$$

Substituting Eq. 46 and Eq. 47 in Eq. 30, the generalized

transmissivity between two blackbodies in the far-field is:

$$\begin{aligned} T_{1 \rightarrow 2}^{bb}(\omega) &= \frac{\omega^2}{c^2} \oint_{S_1} d\mathbf{r} \oint_{S_2} d\tilde{\mathbf{r}} \frac{(-\mathbf{n}_1(\mathbf{r}) \cdot \hat{\mathbf{R}}) (\mathbf{n}_2(\tilde{\mathbf{r}}) \cdot \hat{\mathbf{R}})}{2\pi^2 R^2} \\ &= \frac{\omega^2}{2\pi c^2} A_1 F_{1,2} \end{aligned} \quad (48)$$

and $Q_{1,2}^{bb}$ is given by:

$$\begin{aligned} Q_{1,2}^{bb} &= \int_0^\infty \frac{d\omega}{2\pi} \frac{\omega^2}{2\pi c^2} A_1 F_{1,2} [\Theta(\omega, T_1) - \Theta(\omega, T_2)] \\ &= A_1 F_{1,2} \frac{\pi^2 k_B^4}{60 c^2 \hbar^3} (T_1^4 - T_2^4) \\ &= A_1 F_{1,2} \sigma_{SB} (T_1^4 - T_2^4) \end{aligned} \quad (49)$$

The agreement of obtained results in the examples of Sec. 5.1 with earlier published works [9, 39] and the above derivation for the radiative heat transfer between blackbodies attest to the correctness of our definitions of the generalized transmissivity (Eq. 25, Eq. 27, and Eq. 30). Extension of the proof given here to obtain Eq. 49 to the case of gray body radiative transfer is not a simple one because we consider only specular reflection at surfaces.

6. Summary

In this paper, we have developed a dyadic Green's function formalism to determine radiative heat transfer and non-equilibrium van der Waals/Casimir forces between objects of arbitrary shapes, sizes, and with frequency dependent dielectric permittivity and magnetic permeability. The cross-spectral densities of electromagnetic fields are necessary to evaluate Poynting vector and electromagnetic stress tensor from which radiative transfer and forces between objects can be evaluated. Using Rytov's fluctuational electrodynamics, expressions for cross-spectral densities in terms of volume integrals of products of DGFs are obtained. Green's identities for dyadic functions are then used to convert these volume integral expressions into surface integrals of products of tangential components of the DGFs on the surfaces of scatterers. The spectral radiative transfer between two objects is described in terms of a single quantity - the generalized transmissivity - which can be represented as a double integral of products of tangential components over the surfaces of the two objects. The spectral integral of $T_{1 \rightarrow 2}^e(\omega)$ weighted by the temperature derivative of the Bose-Einstein function yields the thermal radiative conductance between the two objects. In the geometric optics limit, the thermal radiative conductance between two blackbodies, as derived from the generalized transmissivity, is shown to agree with the predictions of the classical theory of radiative transfer.

While many computational methods, such as finite element method, vector eigenfunction expansion, or T-matrix

method, can be employed to compute $T_{1 \rightarrow 2}^e(\omega)$, the surface integral expression in Eq. 30 points towards the surface integral equation method (SIEM) as best-suited for the purpose. The main advantage of a surface integral equation based method is the potential reduction in computational cost due to restriction of the discretization domain to a surface rather than a volume. This advantage becomes more important as the size of the object becomes larger. After the submission of this work, we were made aware of a recently released free software titled SCUFF-EM (**S**urface **C**urrents **F**ield **F**ormulation of **E**lectro-**M**agnetism)¹ based on a surface integral formulation of the electromagnetic scattering problem [40, 41]. The similarities between photon and phonon transport lead us to wonder whether such a surface integral formulation for phonon energy transport in mesoscale structures in the harmonic limit is also possible.

This work has been funded partially by National Science Foundation through Grant CBET-0853723.

References

- [1] Greffet, J.J., Carminati, R., Joulain, K., Mulet, J.P., Mainguy, S., Chen, Y.. Coherent emission of light by thermal sources. *Nature* 2002;416:61.
- [2] Narayanaswamy, A., Chen, G.. Surface modes for near field thermophotovoltaics. *Appl Phys Lett* 2003;82:3544–3546.
- [3] Laroche, M., Carminati, R., Greffet, J.J.. Near-field thermophotovoltaic energy conversion. *J Appl Phys* 2006;100:063704.
- [4] Basu, S., Chen, Y.B., Zhang, Z.M.. Microscale radiation in thermophotovoltaic devices - a review. *Int J Energ Res* 2007;31:689–716.
- [5] Basu, S., Zhang, Z., Fu, C.. Review of near-field thermal radiation and its application to energy conversion. *Int J Energ Res* 2009;33(13):1203–1232.
- [6] Francoeur, M., Menguc, M., Vaillon, R.. Near-field radiative heat transfer enhancement via surface phonon polaritons coupling in thin films. *Appl Phys Lett* 2008;93(4):043109–043109.
- [7] Francoeur, M., Pinar Menguc, M.. Role of fluctuational electrodynamics in near-field radiative heat transfer. *J Quant Spectrosc Radiat Transfer* 2008;109(2):280–293.
- [8] Francoeur, M., Pinar Menguc, M., Vaillon, R.. Solution of near-field thermal radiation in one-dimensional layered media using dyadic Green's functions and the scattering matrix method. *J Quant Spectrosc Radiat Transfer* 2009;110(18):2002–2018.
- [9] Biehs, S.A., Rousseau, E., Greffet, J.J.. Mesoscopic description of radiative heat transfer at the nanoscale. *Phys Rev Lett* 2010;105(23):234301.
- [10] Ben-Abdallah, P., Joulain, K.. Fundamental limits for noncontact transfers between two bodies. *Phys Rev B* 2010;82(12):121419.
- [11] Krüger, M., Emig, T., Kardar, M.. Nonequilibrium electromagnetic fluctuations: Heat transfer and interactions. *Phys Rev Lett* 2011;106(21).
- [12] Narayanaswamy, A., Chen, G.. Thermal near-field radiative transfer between two spheres. *Phys Rev B* 2008;77:075125.
- [13] Sasiithlu, K., Narayanaswamy, A.. Convergence of vector spherical wave expansion method applied to near-field radiative transfer. *Opt Express* 2011;19(104):A772–A785.
- [14] Sasiithlu, K., Narayanaswamy, A.. Proximity effects in radiative heat transfer. *Phys Rev B* 2011;83(16):161406.
- [15] Carrillo, L., Bayazitoglu, Y.. Nanosphere near-field radiative heat-exchange analysis. *J Thermophys Heat Tr* 2010;24(2):309–315.
- [16] Otey, C., Fan, S.. Numerically exact calculation of electromagnetic heat transfer between a dielectric sphere and plate. *Phys Rev B* 2011;84(24):245431.
- [17] Wang, L., Zhang, Z.. Phonon-mediated magnetic polaritons in the infrared region. *Optics Express* 2011;19(102):A126–A135.
- [18] Wang, L., Zhang, Z.. Wavelength-selective and diffuse emitter enhanced by magnetic polaritons for thermophotovoltaics. *Applied Physics Letters* 2012;100(6):063902–063902.
- [19] Zhang, Z., Wang, L.. Measurements and modeling of the spectral and directional radiative properties of micro/nanostructured materials. *International Journal of Thermophysics* 2011;1–34.
- [20] Rodriguez, A., Ilic, O., Bermel, P., Celanovic, I., Joannopoulos, J., Soljačić, M., et al. Frequency-selective near-field radiative heat transfer between photonic crystal slabs: A computational approach for arbitrary geometries and materials. *Phys Rev Lett* 2011;107(11):114302.
- [21] Krüger, M., Bimonte, G., Emig, T., Kardar, M.. Trace formulae for non-equilibrium casimir interactions, heat radiation and heat transfer for arbitrary objects. *arXiv preprint arXiv:12070374* 2012;.
- [22] Landauer, R.. Spatial variation of currents and fields due to localized scatterers in metallic conduction. *IBM J Res Dev* 1957;1(3):223–231.
- [23] Landauer, R.. Conductance determined by transmission: probes and quantised constriction resistance. *Journal of Physics: Condensed Matter* 1989;1:8099.
- [24] Imry, Y., Landauer, R.. Conductance viewed as transmission. *Reviews of Modern Physics* 1999;71(2):306–312.
- [25] Datta, S.. *Electronic transport in mesoscopic systems*. Cambridge Univ Pr; 1997.
- [26] Ben-Abdallah, P., Biehs, S., Joulain, K.. Many-body radiative heat transfer theory. *Phys Rev Lett* 2011;107(11):114301.
- [27] Messina, R., Antezza, M.. Scattering-matrix approach to casimir-lifshitz force and heat transfer out of thermal equilibrium between arbitrary bodies. *Phys Rev A* 2011;84(4):042102.
- [28] Narayanaswamy, A., Chen, G.. Dyadic Green's functions and electromagnetic local density of states. *J Quant Spectrosc Radiat Transfer* 2010;111(12-13):1877–1884.
- [29] Callen, H.B., Welton, T.A.. Irreversibility and generalized noise. *Phys Rev* 1951;83:34–40.
- [30] Landau, L.D., Lifshitz, E.M.. *Statistical Physics*. Addison-Wesley; 1969.
- [31] Eckhardt, W.. First and second fluctuation-dissipation-theorem in electromagnetic fluctuation theory. *Opt Commun* 1982;41(5):305–309.
- [32] Mandel, L., Wolf, E.. *Optical coherence and quantum optics*. Cambridge Univ Pr; 1995.
- [33] Yaghjian, A.D.. Electric dyadic Green's functions in the source region. *Proc IEEE* 1980;68:248–263.
- [34] Tai, C.T.. *Dyadic Green Functions in Electromagnetic Theory*. Piscataway, NJ: IEEE Press; 1993.
- [35] Narayanaswamy, A., Chen, G.. Thermal emission control with one-dimensional metallodielectric photonic crystals. *Phys Rev B* 2004;70:125101.
- [36] Narayanaswamy, A., Chen, G.. *Annual Review of Heat Transfer*; vol. 14; chap. Direct Computation of Thermal Emission from Nanostructures. Begell House; 2005, p. 169–195.
- [37] Wang, L., Basu, S., Zhang, Z.. Direct and indirect methods for calculating thermal emission from layered structures with nonuniform temperatures. *Journal of Heat Transfer* 2011;133:072701.
- [38] Chew, W.C.. *Waves and Fields in Inhomogeneous Media*. Piscataway, NJ: IEEE Press; 1995.
- [39] Antezza, M., Pitaevskii, L., Stringari, S., Svetovoy, V.. Casimir-lifshitz force out of thermal equilibrium. *Phys Rev A* 2008;77(2):022901.
- [40] Reid, M., White, J., Johnson, S.. *Fluctuating surface currents:*

¹It can be downloaded from <http://homerreid.ath.cx/scuff-EM/>

A new algorithm for efficient prediction of Casimir interactions among arbitrary materials in arbitrary geometries. i. theory. arXiv preprint arXiv:12030075 2012;.

- [41] Rodriguez, A., Reid, M., Johnson, S.. Fluctuating surface-current formulation of radiative heat transfer for arbitrary geometries. arXiv preprint arXiv:12061772 2012;.